



INTELLECTUAL PROPERTY OFFICE OF THE PHILIPPINES
 BUREAU OF PATENTS
 Intellectual Property Center
 No. 28 Upper McKinley Road,
 McKinley Hill Town Center,
 Fort Bonifacio, Taguig City
 Tel. No. (02) 238-6300
 Website: <http://www.ipophil.gov.ph>

**SAPALO VELEZ BUNDANG & BULILAN
 LAW OFFICES**
 11TH FLR. SECURITY BANK CENTRE
 6776 AYALA AVENUE, 1200, MAKATI CITY
 METRO MANILA, PHILIPPINES

P13110

IPAS FORM NO.:	
IPAS AGENT CODE:	8002
IPO BOX NO.:	
PAPER NO.:	

SVBB LAW OFFICES
 RECEIVED
 APR 05 2014
 NP4022/ad
 By: Kat

Patent No: Invention 1/2007/502430 Issue Date: 30/01/2014
 Patentee: 4LIFE PATENTS, LLC [US]
 Title: TRANSFER FACTOR PREPARATIONS AND ASSOCIATED METHODS
 International Filing Date 02/05/2006

NOTICE OF ISSUANCE OF LETTERS PATENT

This is to inform you that the Letters Patent Certificate for the above-identified invention has been prepared and is now ready for release.

Said Letters Patent Certificate will be released to you or to your authorized representative upon presentation of this letter to this office.

Please be informed that no Letters Patent Certificate shall be released by the Office unless a documentary stamp(s) worth P20.00 is affixed on the face thereof as required under R.A. 7660 date December 23, 1993 and R.A. 9243 dated February 17, 2004, more particularly Title VII, Sections 173 to 201 thereof.

MAILED

APR 30 2014

THE BUREAU OF PATENTS

Note: this is a system-generated notice. No signature is necessary.

Republic of the Philippines



INTELLECTUAL PROPERTY
PHILIPPINES
BUREAU OF PATENTS

Patent No. 1-2007-502430

Having complied with the provisions of Republic Act No. 8293 and its regulations, this Office grants **LETTERS PATENT** for an


INVENTION

the pertinent data, specification and claim/s of which are hereunto annexed and made part hereof.

Now, therefore, this **LETTERS PATENT** grants unto its owner/s the exclusive right throughout the Philippines to make, use, sell or import the invention and where the invention is or includes a process, including the product obtained directly or indirectly from such process; and,

Unless sooner terminated as provided for by law and the regulations, the term of this **LETTERS PATENT** shall be **TWENTY (20) YEARS** from the date of filing.

IN WITNESS WHEREOF, I have hereunto affixed my hand and the seal of the Intellectual Property Office at Taguig City, Philippines


EPIFANIO M. EVASCO
Director of Patents



**Intellectual Property Office of the Philippines
Bureau of Patents**

[11] Patent No.: 1/2007/502430

[45] Issued: 30/01/2014

[21] Application No: 1/2007/502430
 National Phase Entry Date 05/11/2007
 [22] International Filing Date: 02/05/2006
 [43] International Publication Date: 09/11/2006
 [86] International Application No.: PCT/US2006/017118
 [87] International Publication No.: WO 2006/119408 A1
 [60] Division of Application No.:
 [54] Title: TRANSFER FACTOR PREPARATIONS AND ASSOCIATED METHODS
 [71] Applicant(s): 4LIFE PATENTS, LLC [UTAH, US]
 [72] Inventor(s): CALVIN W. MCCAUSLAND, [UTAH, UNITED STATES OF AMERICA]; BRENT VAUGHAN, [UTAH, UNITED STATES OF AMERICA]; DAVID LISONBEE, [UTAH, UNITED STATES OF AMERICA]; WILLIAM J. HENNEN, [UTAH, UNITED STATES OF AMERICA]
 [30] Foreign Application Priority Data: 60/677,226 02/05/2005 US
 [51] Int. Class⁸: A 61K 35/14, 35/50, 36/00, 36/87
 [58] Field of search: A 61K 35/14, 35/50, 36/00, 36/87
 [56] Reference(s) Considered/Cited:

Category	Document description	Relevant to claim No.	Document No.
	WO2004/017916; 3/4/2004; Hennen, W.		1
	WO2004/041071; 5/21/2004; Dadali et al		2

ABSTRACT

A drink includes an edible liquid or semisolid preparation and transfer factor. The drink may also include lactoferrin and one or more preservatives. An edible preparation includes a fruit component and transfer factor. The fruit component may include at least one oligoproanthocyanidin-containing fruit. The edible preparation may also include lactoferrin. One or more preservatives may also be included in the edible preparation. The drink or the edible preparation may be sterilized or pasteurized.

Industry / Technical Field	Medicinal Preparations			
No. of Claim(s)	27	claim/s	Drawing(s)	0 sheet/s
Multiple Dep. Claim(s)	33	claim/s	Specification and Claim(s)	24 page(s)
Species Claim(s)	0	claim/s	Sequence Listings	0 page(s)
Total No. of Claim(s)	60	claim/s		

U.S. DEPARTMENT OF AGRICULTURE
FOOD SAFETY AND INSPECTION SERVICE
2013 OCT 23 11:41:03

TRANSFER FACTOR PREPARATIONS AND ASSOCIATED METHODS

5

ABSTRACT

A drink includes an edible liquid or semisolid preparation and transfer factor. The drink may also include lactoferrin and one or more preservatives.

10 An edible preparation includes a fruit component and transfer factor. The fruit component may include at least one oligoproanthocyanidin-containing fruit. The edible preparation may also include lactoferrin. One or more preservatives may also be included in the edible preparation. The drink or the edible preparation may be sterilized or pasteurized.

15

20

25

30

35

40

2013 OCT 23 17:14:03

5

TRANSFER FACTOR PREPARATIONS AND ASSOCIATED METHODS

TECHNICAL FIELD

10

The present invention relates generally to preparations that include transfer factor and, more specifically, to edible preparations, such as drinks and solids that include transfer factor. Additionally, the present invention relates to methods for manufacturing edible, transfer factor-containing preparations and to methods that include administration of such preparations.

15

DISCLOSURE OF THE INVENTION

20

In one aspect, the present invention includes a drink that includes transfer factor. The drink includes an edible liquid or semisolid component and transfer factor. The liquid or semisolid component may comprise a fruit juice, a gelatin, a dairy-based product, or any other suitable, drinkable composition with components that are compatible with transfer factor. When mixed with the liquid component, the transfer factor may retain substantially all of one or more of its activities (*e.g.*, components of the liquid component may not interfere with one or more activities of transfer factor), or one or more of the activities of transfer factor may actually be enhanced by one or more components of the liquid component of the drink.

30

In another aspect, the present invention includes a composition with a fruit component and transfer factor. The fruit component may include at least one oligoproanthocyanidin ("OPC")-containing fruit or an extract thereof. The term "extract" is broadly defined herein, including any OPC-including part of a fruit. Examples of extracts include, without limitation, juices (dilute, normal concentration, or concentrate), dehydrated fruit, and powders including one or more components of the fruit. Such a composition may be in

35

a liquid form or a solid form, including but not limited to solid forms that are configured for at least partial dissolution or digestion in the mouth of a subject.

5 Another aspect of the present invention includes a process for making an edible preparation that includes transfer factor. The process includes mixing a fruit component with transfer factor. Preservatives may also be included in the mixture. The mixture may be chilled to prevent microbial growth. To further prevent microbial growth, the mixture may be pasteurized
10 before chilling. Alternatively, the mixture may be sterilized.

Other features and advantages of the present invention will become apparent to those of ordinary skill in the art through consideration of the ensuing description and the appended claims.

15

BEST MODES FOR CARRYING OUT THE INVENTION

An exemplary embodiment of a drink that includes transfer factor may be liquid or semisolid. Such a drink may include an edible liquid or semisolid
20 component, as well as transfer factor. The liquid or semisolid component may, by way of nonlimiting example, comprise a fruit juice, a gelatin, a dairy-based product, or any other suitable, drinkable composition with components that are compatible with transfer factor. When mixed with the liquid component, the transfer factor may retain substantially all of one or
25 more of its activities (*e.g.*, components of the liquid component may not interfere with one or more activities of transfer factor), or one or more of the activities of transfer factor may actually be enhanced by one or more components of the liquid component of the drink. Such a drink may also include one or more preservatives. Alternatively, or in addition, lactoferrin
30 may be included in a drink that includes transfer factor.

Another embodiment of an edible preparation that includes transfer factor also includes a fruit component. The edible preparation may also

include one or more preservatives. Alternatively, or in addition, lactoferrin may be included in such an edible composition.

5 The fruit component includes at least one fruit that naturally includes OPC or a juice or other extract of such a fruit. By way of nonlimiting example, the fruit component may include one or more of açai, elderberry, grape, and pomegranate or an extract thereof. OPC is a known antioxidant and may, therefore, be useful in neutralizing or otherwise acting against free radicals and other oxidants, which may adversely affect cell membranes, cause accelerated cellular aging, and are known or believed to be at least indirectly responsible for a wide variety of disease states, as well as compromised immunity, in living beings.

15 The transfer factor component may include any type of transfer factor, as well as a combination of two or more types of transfer factor. For example, avian transfer factor, bovine transfer factor, or any other type of transfer factor may be included in the transfer factor component. The transfer factor of the transfer component factor may be derived from any suitable, acceptable source. For example, avian transfer factor may be obtained from eggs, such as by a process disclosed in U.S. Patent 6,468,534 to Hennen et al. (hereinafter "Hennen"). An example of the manner in which bovine transfer factor may be obtained is disclosed in U.S. Patent 4,816,563 to Wilson et al. (hereinafter "Wilson"). Compositions that include two or more types of transfer factor, as well as processes for combining and processing two or more types of transfer factor, are disclosed in U.S. Patent 6,866,868 to Lisonbee et al. (hereinafter "Lisonbee").

30 Transfer factor is known or believed to improve the oxidative balance of a living being, as well as to enhance the effectiveness of antioxidants, as demonstrated by the disclosure of the international patent application filed pursuant to the Patent Cooperation Treaty and having International Publication Number WO 2004/041071 A2 (hereinafter "Dadali").

An edible preparation according to the present invention may also include one or more preservatives. Suitable preservatives, such as those accepted for use in foods and beverages, may be used. Examples of preservatives that may be included in an edible preparation of the present invention include, but are not limited to, sodium benzoate and preservatives from the paraben family of chemicals.

Lysozyme, when used in an edible preparation that incorporates teachings of the present invention, acts as a preservative. While lysozyme has been used as a preservative in cheeses, it is not believed to have been previously used in this capacity in edible preparations that include fruits or extracts thereof.

Lactoperoxidase may also, or alternatively, be included in edible preparations that incorporate teachings of the present invention. Lactoperoxidase is another preservative that has been used in dairy products, but it is believed that lactoperoxidase has not been used to preserve edible compositions that include fruits or extracts of fruits.

Lactoferrin is known or believed to stimulate the immune system and may work in concert with transfer factor to improve the immunity of a subject that receives an edible preparation that incorporates teachings of the present invention. Lactoferrin is also known to starve bacteria and, thus, may act as a preservative when included in an edible preparation that incorporates teachings of the present invention.

In an exemplary embodiment, the edible preparation may be a liquid, a semisolid, or a solid. A liquid form of edible preparation may be in the form of a fruit juice. A solid form of edible preparation may be configured for consumption as a solid (*e.g.*, as a chewable tablet, an effervescent tablet, a dissolvable wafer, a dissolvable gel strip, etc.), for reconstitution as a liquid, or in any other suitable fashion.

A liquid edible preparation that incorporates teachings of the present invention is described in the following example:

EXAMPLE 1

5

An example of a formulation for a liquid preparation that includes transfer factor follows:

TABLE 1

10

Ingredient	% of Total (w/v)	Density (g/ml)	% of Total Juices (v/v)
Water	76.681	1.000	
Apple Juice	5.146	1.346	19
Purple Grape Juice	5.100	1.330	19
Glycerine	3.980	1.249	
Blueberry Juice	3.772	1.315	18
Transfer Factor E-XF	1.912		
Pomegranate Juice	1.480	1.315	15
Grape color concentrate (e.g., MEGANATURAL™ purple from Canandaigua Concentrates & Colors, a Division of Canandaigua Wine Company of Madera, California)	0.500	1.306	
Vitamin C	0.498		
Elderberry Juice	0.419	1.315	15
Flavorings			
Berry flavor (BE-01407)	0.198	1.000	
Berry flavor (BE-01271)	0.055	1.000	
Natural Vanilla (VA-01239)	0.165	1.000	
Açai Powder	0.318		14
Lactoferrin	0.191		
Lysozyme	0.014		
Lactoperoxidase	0.003		

Transfer Factor E-XF includes bovine transfer factor from cow colostrum and avian transfer factor from the yolk of a chicken's egg.

15

The flavorings listed in TABLE 1 are available from Flavors Inc.

A daily dosage of about one fluid ounce (about 30 ml) or more of a composition with ingredients in the proportions listed in TABLE 1 may be administered to or consumed by a subject. In addition to the numerous known and believed benefits of antioxidants, including the benefits of OPC
5 and OPC-containing fruits such as açai, administration or consumption of an edible composition that incorporates teachings of the present invention provides the subject with the additional and sometimes synergistic beneficial affects of transfer factor, which are known in the art, as evidenced by the disclosures of Dadali, Hennen, Lisonbee, and Wilson.

10

An edible preparation may be made by mixing components of a food base with transfer factor by processes that are known in the art. Additionally, lactoferrin or preservatives, including without limitation lysozyme, lactoperoxidase, and other food preservatives, may be mixed with the food
15 base.

Of course, the types of processes that are used and, possibly, the order in which certain ingredients are included, may depend in part upon the form, or state (*e.g.*, liquid, solid, semisolid, etc.), of the various ingredients that are
20 mixed with one another, their solubilities, and the desired form, or state, of the resulting edible preparation (*i.e.*, whether the edible preparation is liquid, semisolid, solid, includes carbonation, etc.). Suitable processes that may be used to manufacturing edible preparations of a variety of different forms are well known and within the skill of those in the relevant art.

25

A liquid edible preparation may be manufactured with liquid ingredients or liquid and dissolvable solid (*e.g.*, powder, crystalline, etc.) or semisolid (*e.g.*, gel, paste, etc.) ingredients. Alternatively, dry ingredients may be mixed with one another, then reconstituted (*e.g.*, in water, juice, etc.)
30 by either the manufacturer, a distributor, or an end-user to liquid form.

Known techniques, such as those disclosed in "Principles and Practices of Small- and Medium-Scale Fruit Juice Processing," Food and

Agricultural Organization of the United Nations (FAO) Services Bulletin 146 (Rome, 2001), may be used in one or more parts of a process for manufacturing fluid edible preparations that incorporate teachings of the present invention.

5

A solid or semisolid edible preparation may be manufactured with dry, semisolid, or liquid components, or combinations thereof, then, if desired, dried (*e.g.*, by dehydration processes, etc.) to a desired state. A solid edible preparation may be manufactured in any suitable form (*e.g.*, chewable
10 tablet, dissolvable wafer or gel, chewing gum, reconstitutable powder, etc.) by processes that are well known in the art. Of course, various additional ingredients (*e.g.*, fillers, sweeteners, wicking agents, etc.) may be used to facilitate manufacture of the edible preparation in a desired solid form. Examples of additional ingredients that may be included in a solid or
15 semisolid edible preparation include, but are not limited to, one or more of the following: pullulan, corn starch, gelatin, dextrin, glycerin, carrageenan, xanthan gum, dextrose, corn syrup, and bees wax. The incorporation of additional ingredients into an edible composition is well within the skill of one who practices in the art.

20

Without limiting the scope of the present invention, various types of solid edible compositions may be formed by known processes, including, but not limited to, the processes that are disclosed at "Tablets,"
<http://pharmlabs.unc.edu/tablets/text.htm> and in U.S. Patents 6,326,028 to
25 Nivaggioli et al., 6,733,781 to Abu-Izza et al., and 6,811,795 to Wehling.

Processes that are used to manufacture edible preparations that are in a form that is not completely dry, such as a liquid or semisolid state, may be effected at a low temperature (*e.g.*, between about 0° C. and about 10° C., at
30 about 4° C, etc.), such as in a refrigerated environment, then transported and stored at such temperatures to reduce the likelihood of microbial growth or proliferation therein.

Alternatively, an edible preparation that is in a form that is not completely dry may be pasteurized or sterilized. Pasteurization processes, which decrease the number of microorganisms present, but do not entirely eliminate the microorganisms, improve the stability of products that are to be stored at reduced temperatures (*e.g.*, frozen or refrigerated, or "chilled").
5 When an edible preparation is sterilized, all or substantially all microorganisms therein are killed or inactivated, facilitating prolonged storage of the edible preparation at room temperature or even higher temperatures.

10

As an example, an edible preparation that includes transfer factor may be sterilized by known superheated steam injection processes. The temperatures and durations of such processes depend, of course, upon the form and ingredients of the composition to be sterilized. When making a liquid preparation, the resulting edible preparation may be "flash" heated to a particular temperature (*e.g.*, 120° C.) for a corresponding duration (*e.g.*, two seconds). Alternatively, a sterilization or pasteurization process of different duration and temperature may be used, so long as the duration and temperature of the process are in substantial accord with a practice that has
15 been accepted in the art, such as use of the following equation:
20

$$t_p = 5 \cdot 10^{14} \cdot e^{-0.4353 \cdot T_{mo}},$$

where t_p is the minimum duration of the process, and T_{mo} is the temperature at which the process is effected.
25

Of course, processes that reduce microbial load on an edible preparation of the present invention need not comprise heat-treatment techniques. Sterilization or other microbial load-reducing techniques that employ other means (*e.g.*, filtration, antimicrobial ingredients, etc.) may also
30 be used in manufacturing an edible preparation. Examples of suitable processes are disclosed in Hughes, D. E., and Nyborg, W., "Minimally

Processed Fruits and Vegetables: Reducing Microbial Load by Nonthermal Physical Treatments," Food Technology 52(6): 66-71 (1997).

5 It is desirable that, following pasteurization or sterilization, the transfer factor retain some if not substantially all or all of its activity. A variety of pasteurization or sterilization processes may be employed, including pasteurization or sterilization processes that may be used to reduce microbial counts or completely eliminate microorganisms from foods. As many sterilization processes are known to significantly reduce the activity of
10 certain proteins, including antibodies, a study was performed to determine whether transfer factor retains at least some of its activity following sterilization.

In the study, mouse footpad assay techniques, similar to those
15 disclosed in J. Natl. Cancer Inst. 55(5):1089-95 (Nov. 1975), were used to determine the affects of heat pasteurization or sterilization processes (specifically, superheated steam injection processes) on edible preparations including transfer factor. Two sterilized samples were compared with an unsterilized sample, as well as with a negative control and a positive control.

20 Separate populations of six mice were tested for each of the five samples and controls. The tests were conducted in two phases, a first that immediately followed heat sterilization of the samples, and a second that was conducted after storing the two heat sterilized samples at a temperature of
25 about 40° C. for about three months, which is well-accepted in the art to be the equivalent of about one year of storage at room temperature. Thirty different mice were used in each phase of the study. The following procedures were followed in each phase of the study

30 In the positive control (*i.e.*, the "fifth group"), fourteen days prior to testing, the footpads of the right rear feet of six BALB/c mice having ages of about nine weeks to about ten weeks were anesthetized with isoflurane. Then 0.02 ml of an about 50/50 (wt/wt) mixture of Freund's adjuvant and bovine

rhinotracheitis virus diarrhea vaccine was administered intramuscularly to each mouse by way of two injections at the base of each side of the mouse's tail. This early injection of antigen allows the mice of the positive control group to elicit their own primary immune response and secondary, or
5 delayed-type hypersensitivity response to the antigen. The mice of the other five groups were not preexposed to the antigen in this manner.

About twenty-four hours before evaluating the hind footpads of the mice, the six BALB/c mice of each group, which were of similar age to the
10 mice of the positive control group, were anesthetized with isoflurane. About 0.5 ml of a sample solution or control solution was then administered by subcutaneous injection at the back of the neck of each mouse.

In the first group (*see* EXAMPLE 2 below), which was the negative
15 control group, the back of the neck of each mouse was injected with about 0.5 ml of sterile saline solution.

In the second group (*see* EXAMPLE 3 below), the sample solution included 16% solids (w/v) of a reconstituted (in distilled, deionized water)
20 lyophilized colostrum fraction that included transfer factor. The solution was set at a pH of 4.0, which was intended to estimate the pH of a fruit juice preparation (the actual pH of which is about 3.6 or about 3.7). Following reconstitution and pH adjustment, the solution was sterilized by heating the same to a temperature of about 120° C. for about two seconds.

25 In the third group (*see* EXAMPLE 4 below), the sample solution included 16% solids (w/v) of a reconstituted (in distilled, deionized water) lyophilized colostrum fraction that included transfer factor. The pH of the resulting solution was not adjusted and, thus, was neutral (*i.e.*, 7.0) or slightly
30 basic (*i.e.*, greater than 7.0)). Following reconstitution, the solution was sterilized by heating the same to a temperature of about 120° C. for about two seconds.

In the fourth group (*see* EXAMPLE 5 below), the sample solution was a concentrate of a colostrum fraction that included transfer factor, which had been diluted to about 16% solids (w/v) in distilled, deionized water. This solution was not heat sterilized or pH adjusted.

5

The mice of the fifth group (*see* EXAMPLE 6 below), which was the positive control groups, respectively, received sterile saline solution.

At the start of the mouse footpad assay, the right hind footpad and the left hind footpad of each mouse were measured, such as with a Starrett gauge. The right hind footpad of each of the thirty mice during each phase of the study was then subcutaneously injected with an antigen-containing solution. The footpad on the left hind foot of each of the thirty mice in each phase, which was used as a control, was injected with about the same volume of a control solution, such as a sterile saline diluent, as the volume of antigen-containing solution that was injected into right hind footpad.

10
15

After a sufficient amount of time (*e.g.*, about twenty-four hours) for the secondary immune response components of the immune system of each mouse to respond, each mouse was again anesthetized and the distances across right and left hind footpads were again measured. A significant amount of swelling, determined by an increase in the distance across a right hind footpad of a mouse from the initial measurement to the second measurement, is indicative of the occurrence of a delayed-type hypersensitivity reaction in that footpad.

20

25

The results of the mouse foot pad assays, and some accompanying analysis, are set forth in EXAMPLES 2 through 5 and 7:

30

EXAMPLE 2

In the first phase of the study, the footpads on the right hind feet of the six mice of the negative control, or first group, exhibited, on average, about

6.35 micrometers more swelling about twenty-four hours after they were injected with the antigen solution than the swelling measured in the footpads of the left hind feet of these mice, which were merely inoculated with sterile saline.

5

The results for the negative control group during the second phase of the study are set forth in the following table:

TABLE 2

10

Mouse	Foot (left/right)	Foot Pad (untreated) (micrometers)	Foot Pad (final) (micrometers)	Foot Pad (difference) (micrometers)
1	Left (control)	1930.40	1955.80	25.40
	Right (test)	1905.00	1930.40	25.40
2	Left (control)	1981.20	2006.60	25.40
	Right (test)	2006.60	2057.40	50.80
3	Left (control)	2057.40	2057.40	0.00
	Right (test)	2032.00	2057.40	25.40
4	Left (control)	2006.60	2032.00	25.40
	Right (test)	2032.00	2057.40	25.40
5	Left (control)	1955.80	2006.60	50.80
	Right (test)	1930.40	1955.80	25.40
6	Left (control)	1905.00	1930.40	25.40
	Right (test)	1876.60	1955.80	76.20

15

Similar to the results from the first phase, the footpads of the right hind feet of the mice of the negative control group exhibited, on average, only 12.70 micrometers more swelling about twenty-four hours after antigen injection than the footpads of the left hind feet of the same mice exhibited twenty-four hours after sterile saline injection. As twenty-four hours is not a sufficient period of time for a mouse to mount a primary (*i.e.*, antibody-mediated) immune response to the antigen, these insignificant differences in swelling

show that the mice did not exhibit a significant secondary immune response to the antigen.

EXAMPLE 3

5

In the first phase of the study, about twenty-four hours after they were injected with the antigen solution, the footpads on the right hind feet of the six mice of the second group of mice (which mice had previously been inoculated with a solution including 16% solids (w/v) colostrum at pH=4.0) swelled, on average, by 50.80 micrometers more than the swelling that was measured in the footpads of the left hind feet of these mice. These results indicate that there was a greater secondary, or delayed-type hypersensitivity, immune response in the footpads into which antigen was injected than in the footpads into which no antigen was injected, which were likely swollen merely because they were pierced by a needle.

15

In the second phase of the study, similar results were obtained, as set forth in the following table:

20

TABLE 3

Mouse	Foot (left/right)	Foot Pad (untreated) (micrometers)	Foot Pad (final) (micrometers)	Foot Pad (difference) (micrometers)
1	Left (control)	1955.80	2006.60	50.80
	Right (test)	1981.20	2057.40	76.20
2	Left (control)	1930.40	2006.60	76.20
	Right (test)	1955.80	2108.20	152.40
3	Left (control)	1955.80	2006.60	50.80
	Right (test)	1981.20	2082.80	101.60
4	Left (control)	2032.00	2057.40	25.40
	Right (test)	2057.40	2108.20	50.80
5	Left (control)	1930.40	2006.60	76.20

	Right (test)	1955.80	2032.00	76.20
6	Left (control)	2057.40	2108.20	50.80
	Right (test)	2032.00	2159.00	127.00

More specifically, the footpads of the right hind feet of the six mice of the second group swelled so that they measured, on average, 42.33 micrometers more than the swelling that was measured in the footpads of the left hind feet of these mice before and after inoculation of their foot pads with the antigen solution. The similar results between the first and second phases of the study indicate that, once a liquid solution that includes transfer factor has been heat sterilized, there is little or no change in the activity of the transfer factor after prolonged storage of the solution.

10

EXAMPLE 4

The results for the third group of mice (which mice had previously been inoculated with a solution including 16% solids (w/v) colostrum at normal pH) were similar to the results for the second group in the first and second phases of the study.

15

In the first phase of the study, about twenty-four hours after the footpad injections, the antigen solution-inoculated footpads on the right hind feet of the six mice of the third group of mice swelled, on average, by 35.98 micrometers more than the swelling that was measured in the sterile saline-inoculated footpads of the left hind feet of these mice. These results indicate that there was a greater secondary, or delayed-type hypersensitivity, immune response in the footpads into which antigen was injected than in the footpads into which no antigen was injected, which were likely swollen merely because they were pierced by a needle.

25

In the second phase of the study, similar results were obtained, as set forth in the following table:

TABLE 4

Mouse	Foot (left/right)	Foot Pad (untreated) (micrometers)	Foot Pad (final) (micrometers)	Foot Pad (difference) (micrometers)
1	Left (control)	2006.60	2032.00	25.40
	Right (test)	2032.00	2082.80	50.80
2	Left (control)	2057.40	2057.40	0.00
	Right (test)	2006.60	2108.20	101.60
3	Left (control)	1981.20	2006.60	25.40
	Right (test)	2057.40	2082.80	25.40
4	Left (control)	2006.60	2057.40	50.80
	Right (test)	2032.00	2082.80	50.80
5	Left (control)	2057.40	2082.80	25.40
	Right (test)	2082.80	2159.00	76.20
6	Left (control)	2082.80	2108.20	25.40
	Right (test)	2108.20	2159.00	50.80

5 These results show that the footpads of the right hind feet of the six mice of the third group swelled so that they measured, on average, 33.87 micrometers more than the swelling that was measured in the footpads of the left hind feet of these mice before and after inoculation of the foot pads with the antigen solution. The similar results between the first and second phases of the study indicate that, following prolonged storage, there was little or no change in the activity of the transfer factor in a heat-sterilized solution.

10

EXAMPLE 5

15 These results were confirmed by the results that were obtained from the fourth group of mice. In particular, during the first phase of the study, the footpads of the right hind feet of mice in the fourth group (which included mice that had been inoculated with a diluted liquid colostrum fraction that was not heat sterilized) exhibited, on average, about 35.98 micrometers more swelling than the foot pads of left hind feet of these mice about twenty-four

EXAMPLE 6

This conclusion was verified by data from another mouse footpad assay, in which six BALB/c mice were inoculated, behind the neck, with 0.5 ml of a solution including 16% solids (w/v) of a spray-dried colostrum fraction that had been reconstituted in distilled, deionized water. About twenty-four hours later, the mice were anesthetized with isoflurane, then footpads on their hind feet measured and inoculated in the manner described above (*i.e.*, left footpad with sterile saline, right footpad with the antigen solution). After about another twenty-four hours, the footpads were again measured. The right footpads of these mice swelled, on average, about 42.33 micrometers more than the footpads on the left hind feet of these mice. This value is comparable to (*i.e.*, not significantly different from) the differences noted above with respect to the second, third, and fourth groups of mice in both the first and second phases of the study detailed in EXAMPLES 2 through 5 and 7, further supporting the conclusion that heat sterilization of a solution that includes transfer factor, such as the solutions that were tested on the second and third groups of mice (EXAMPLES 3 and 4) does not have a significant adverse affect on the activity of the transfer factor.

20

EXAMPLE 7

The fact that the transfer factor with which the mice were inoculated was responsible for the increased secondary immune response is supported by the results from the fifth group, or positive control group, of mice during the second phase of the study, as set forth in the following table:

TABLE 6

Mouse	Foot (left/right)	Foot Pad (untreated) (micrometers)	Foot Pad (final) (micrometers)	Foot Pad (difference) (micrometers)
1	Left (control)	1981.20	2006.60	25.40

	Right (test)	2006.60	2082.80	76.20
2	Left (control)	1828.80	1854.20	25.40
	Right (test)	1879.60	2082.80	203.20
3	Left (control)	1905.00	1930.40	25.40
	Right (test)	1981.20	2082.80	101.60
4	Left (control)	2006.60	2057.40	50.80
	Right (test)	2032.00	2184.40	152.40
5	Left (control)	2032.00	2057.40	25.40
	Right (test)	2057.40	2184.40	127.00
6	Left (control)	2108.20	2108.20	0.00
	Right (test)	2082.80	2184.40	101.60

These results, which show on average, 101.60 micrometers more swelling in the footpads that were inoculated with antigen solution over those that were inoculated with sterile saline, are similar to the 124.88 micrometer difference
 5 seen in the mice of the positive control group during the first phase of the mouse footpad study. The greater swelling in the antigen solution-inoculated footpads of the mice of the positive control group is indicative of a greater secondary immune response than that induced artificially by administration of transfer factor, as the mice of the positive control group had a sufficient
 10 period of time (*i.e.*, two weeks) to generate their own transfer factor and, thus, to mount their own secondary immune response to the antigen.

Once an edible preparation of the present invention has been manufactured, it may be introduced into a clean or sterile container for
 15 subsequent transport and storage.

EXAMPLE 8

In another study, mouse footpad assays were conducted to determine
 20 the effectiveness of transfer factor in heat-treated samples of a liquid solution that included transfer factor that had been stored for one year. In total, four samples were prepared, two each having a pH of about 4 and two each having

a pH of about 7. All of the samples had been flash sterilized at a temperature of about 120° C. for about two seconds to about four seconds. The samples were subsequently stored for one year, with one each of the pH = 4 and pH = 7 samples having been stored at room temperature (which varied from about 5 18° C. to about 25° C.) and one each of the pH = 4 and pH =7 samples having been refrigerated (at temperatures of about 4° C.). After one year, the samples were lyophilized. Prior to testing, the lyophilized samples were reconstituted to desired concentrations, then administered in the manner described above.

10

In a first sample, which included liquid having a pH of about 4 that was stored at room temperature, footpad swelling was, on average, 50.80 micrometers greater in footpads that had been injected with antigen versus footpads that had merely been injected with saline. These results were 15 repeated in second (liquid of a pH of about 7 that was stored at room temperature), third (liquid of a pH of about 4 that was refrigerated), and fourth (liquid of a pH of about 7 that was refrigerated) samples, in which hind footpads that had been injected with antigen were, on average, respectively swollen 59.27, 67.73, and 63.50 micrometers more than hind footpads that 20 were merely injected with saline.

Additionally, positive and negative controls were prepared as discussed above. In the positive control, the average difference in swelling between antigen-injected footpads and saline-injected footpads was 114.30 25 micrometers. In the negative control, the average difference in swelling between antigen-injected footpads and saline-injected footpads was only 38.10 micrometers.

Taken together, these data indicate that the increased swelling was due 30 to the presence of transfer factor in the mice in the areas (hind footpads) into which antigen was introduced. Additionally, these data indicate that the transfer factor lost little or none of its effectiveness after heat-treatment and prolonged storage. The activity of transfer factor in refrigerated samples

2013 OCT 23 P/i 4/d/g

CLAIMS

What is claimed:

1. A nutritional supplement, comprising:
5 a fruit component including an oligoproanthocyanidin-containing fruit extract; and
a transfer factor component including transfer factor.

- 10 2. The nutritional supplement of claim 1, in a liquid form.

3. The nutritional supplement of claim 2, wherein the liquid comprises fruit juice.

- 15 4. The nutritional supplement of claim 1, in a solid form.

5. The nutritional supplement of claim 4, further comprising:
a chewable base.

- 20 6. The nutritional supplement of any one of claims 1-5, wherein the oligoproanthocyanidin-containing fruit comprises açai, elderberry, grape, or pomegranate.

- 25 7. The nutritional supplement of any one of claims 1-5, wherein the transfer factor component comprises avian transfer factor or bovine transfer factor.

- 30 8. The nutritional supplement of any one of claims 1-5, further comprising:
at least one preservative.

9. The nutritional supplement of claim 8, wherein the at least one preservative comprises at least one of lysozyme and lactoperoxidase.

10. The nutritional supplement of any one of claims 1-5, further comprising:
lactoferrin.
- 5 11. The nutritional supplement of claim 1, wherein the fruit containing component and the transfer factor component are pasteurized.
12. A method for manufacturing a nutritional supplement, comprising:
10 forming a mixture including a transfer factor component and a fruit component that includes an oligoproanthocyanidin-containing fruit extract.
13. The method of claim 12, further comprising:
15 sterilizing the mixture.
14. The method of claim 12, further comprising:
pasteurizing the mixture.
- 20 15. The method of any of claims 12-14, further comprising:
chilling the mixture.
16. The method of claim 12, wherein forming the mixture comprises forming a mixture further including at least one preservative.
25
17. The method of claim 12, wherein forming the mixture comprises forming a mixture including lactoferrin, lysozyme or lactoperoxidase.
- 30 18. The method of any one of claims 12-14, wherein forming the mixture comprises forming a mixture including a liquid component.

19. The method of claim 18, wherein forming the mixture comprises forming a mixture including an oligoproanthocyanidin-containing fruit juice.

5 20. The method of claim 18, wherein forming the mixture comprises forming a mixture including the oligoproanthocyanidin-containing fruit extract in a dry form.

21. The method of claim 18, further comprising:
10 drying the mixture.

22. The method of claim 12, wherein forming the mixture comprises forming a mixture including only dry components.

15 23. An edible fluid, comprising:
a liquid or semisolid edible preparation including an oligoproanthocyanidin-containing fruit or an oligoproanthocyanidin-containing fruit extract; and transfer factor.

20 24. The edible fluid of claim 23, wherein the oligoproanthocyanidin-containing fruit or the oligoproanthocyanidin-containing fruit extract comprises açai or an açai extract.

25 25. The edible fluid of claim 23 or claim 24, wherein the transfer factor comprises avian transfer factor.

26. The edible fluid of claim 23 or claim 24, wherein the transfer
30 factor comprises bovine transfer factor.

27. The edible fluid of claim 26, wherein the transfer factor further comprises avian transfer factor.