4LIFE TRANSFER FACTOR® CHEWABLE

90 CT/BOTTLE
ITEM CODE #: 170024042
SUGGESTED RETAIL PRICE:
PHP 3,900 | 35 LP

CHEW-LY AMAZING!

4Life
TOGETHER, BUILDING PEOPLE®

MAHALAGANG PAALALA: ANG MGA PRODUKTO NG 4LIFE AY HINDI GAMOT AT HINDI DAPAT GAMITIN PANGGAMOT SA ANUMANG URI NG SAKIT.
IMPORTANT REMINDER: 4LIFE PRODUCTS ARE NOT MEDICINE AND CANNOT BE USED TO CURE ANY DISEASES.
*THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THESE PRODUCTS ARE NOT INTENDED TO DIAGNOSE, TREAT, CURE, OR PREVENT ANY DISEASE.

Nutrition Information

Serving Size: One (1) Tablet
Servings Per Container: 90

	Amount Per Serving
Calories 4 kcal	Calories from Fat 0.3 kcal
Total Fat	0.04 g
Saturated Fat	0.02 g
Trans Fat	0 g
Cholesterol	2 mg
Sodium	1 mg
Total Carbohydrates	0.8 g
Dietary Fiber	0.02 g
Sugar	0.7 g
Total Protein	0.1 g
4Life® Tri-Factor® Formula	200 mg

PRIMARY SUPPORT:
Immune System, Overall Wellness*

INGREDIENTS:
Transfer Factor - tiny messenger molecules composed of amino acids (i.e., Transfer Factors are a unique category of bioactive peptide), that transfer immune memory from an experienced immune system "donor" to a less experienced recipient immune system.*

DID YOU KNOW?

4LIFE TRANSFER FACTOR®

DISCOVERY

- Dr. H. Sherwood Lawrence discovered transfer factor molecules in 1949.
- 4Life secured patents (U.S.# 6,866,868 B1 and U.S.# 7,815,943 B1) that protect the process and use of Transfer Factors derived from colostrum ("first milk") and chicken egg yolks.

BENEFITS

- Transfer Factors support the immune system's natural ability to recognize, respond to, and remember potential threats.*
- Transfer Factors may boost or balance the immune system response to a potential threat.*
- 4Life Transfer Factor Tri-Factor Formula promotes healthy T-cell and B-cell function.*

Test results obtained from an independent, unpublished *in vitro* experiment conducted within the Blokhin Cancer Research Center, at the Russian Academy of Medical Sciences, in Kashirskoe Shosse, Russia. The randomized and controlled *in vitro* study assessed the effects of 4Life Transfer Factor® Classic, 4Life Transfer Factor® Tri-Factor® Formula, or 4Life® Transfer Factor PLUS® Tri-Factor® Formula, versus a positive control (Interleukin-2, or IL-2) on NK cell activity and effectiveness in destroying damaging cells. Blood was collected from healthy volunteers and then incubated for up to 48 hours. (REFERENCE: Kisielevsky MV & Khalturina EO. Antitumor and cytotoxic activity of mononuclear blood cells. *Unpublished observations*)