4Life Transfer Factor Classic Raise Your Immune I.Q.

- Supports healthy immune system function*
- Educates immune cells*

What is 4Life Transfer Factor Classic?

Raise Your Immune I.Q. with 4Life Transfer Factor Classic! This product supports immune system function with UltraFactor XF®, a concentrate of transfer factor molecules from cow colostrum. These immune messenger molecules support the immune system's ability to more effectively recognize, respond to, and remember potential health threats.*

Key features

- Supports healthy immune system function that, in turn, promotes healthy energy levels and the healthy function of all other systems throughout the body*
- Contains transfer factors (immune messenger molecules) and other natural components from cow colostrum that help educate immune cells*
- Increases natural killer (NK) cell function*[†]



DIRECTIONS: Take three (3) capsules daily with 8 oz of fluid.

Serving Size: Three (3) Capsules Servings Per Container: 30 Amount Per Serving UltraFactor XF® A proprietary concentrate of ultra-filtered 4Life Transfer Factor® proteins and other peptides from cow colostrum.

OTHER INGREDIENTS: Gelatin capsule and maltodextrin. CONTAINS INGREDIENTS FROM MILK.

PRIMARY SUPPORT:

Immune System*
Overall Wellness*

SECONDARY SUPPORT:

Heart Health* Energy*

Did you know?

4Life Transfer Factor Classic is 4Life's original formula, containing transfer factors from cow colostrum. Since this product was first released in 1998, 4Life has pioneered Transferceutical® Science, introducing more than 25 new products that contain transfer factors.

Ordering Information

Item # 24080 - 90 ct/bottle Item # 24081 - 12 for the price of 10

†Test results obtained from an independent, unpublished in vitro experiment conducted at the Russian Academy of Medical Sciences, in Kashirskoe Shosse, Russia. The randomized and controlled in vitro study assessed the effects of 4Life Transfer Factor® Classic, 4Life Transfer Factor® Formula, or 4Life® Transfer Factor® Formula, versus a positive control (Interleukin-2, or IL-2) on NK cell activity and effectiveness in destroying damaging cells. Blood was collected from healthy volunteers and then incubated for up to 48 hours. (REFERENCE: Kisielevsky MV & Khalturina EO. Unpublished observations)

*THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE, OR PREVENT ANY DISEASE.

